

Evaluation of the Reach and Impact of the *100% Jeune* Youth Social Marketing Program in Cameroon: Findings from Three Cross-Sectional Surveys

Andrea Plautz¹
Dominique Meekers²

Corresponding author: Dr. Dominique Meekers

¹ Population Services International (PSI), 1120 19th Street, Suite 600, Washington, DC, 20036, aplautz@psi.org

² Department of International Health and Development, Tulane University School of Public Health and Tropical Medicine, 1440 Canal Street, Suite 2200, New Orleans, LA 70131, dmeekers@tulane.edu

Evaluation of the Reach and Impact of the *100% Jeune* Youth Social Marketing Program in Cameroon: Findings from Three Cross-Sectional Surveys

Abstract

Background: The *100% Jeune* youth social marketing program in Cameroon aims to address the high STI/HIV prevalence rates and the high levels of unwanted pregnancy. This study evaluates the *100% Jeune* program, analyzing its reach and impact on condom use, level of sexual activity, and predictors of condom use.

Methods: This analysis uses data from three waves of the Cameroon Adolescent Reproductive Health Survey, implemented at 18-month intervals between 2000 and 2003. The sample is restricted to unmarried youth aged 15-24; sample sizes are 1,956 youth in 2000, 3,237 in 2002, and 3,370 in 2003. Logistic regression analyses determine trends in reproductive health behavior and their predictors, as well as estimate the effect of program exposure on these variables. All regression analyses control for differences in sample characteristics.

Results: A comparison of trends over the 36-month study period shows that substantial positive changes occurred among youth. Results of dose response analyses indicate that some of these positive changes in condom use and predictors of use can be attributed to the *100% Jeune* youth social marketing program. The program contributed to substantial increases in condom use, including consistent use with regular partners among youth of both sexes. Among males, it also contributed to consistent use with casual partners. The program did not decrease the level of sexual activity or reduce the number of sexual partners, despite efforts to promote abstinence. Observed secular trends indicate that factors besides the *100% Jeune* program also contributed to the improvement observed in several areas.

Conclusions: Results show that *100% Jeune* successfully used a variety of mass media and interpersonal communication channels to reach a high proportion of youth throughout the intervention period. In a context in which a variety of governmental and nongovernmental partners are increasing youth-focused reproductive health programming, the *100% Jeune* program reached a higher proportion of youth than did other programs. Collective efforts of multiple organizations over time can lead to improvements in adolescent reproductive health. Resources should be allocated to identify and understand predictors of abstinence and partner reduction to inform future programming decisions.

Introduction

Human immunodeficiency virus (HIV) rates in Cameroon have increased steadily in the past decade. Although the national HIV prevalence was only 2.9% in 1994, it is now estimated at close to 12% (Matagne et al., 1998; Mbopi et al., 1998; UNAIDS, 2002; UNICEF, 2002). There is evidence that youth in Cameroon are among the populations most at risk for contracting HIV. In 2002, UNAIDS reported an HIV prevalence of 11.5% among pregnant 15–19-year-olds and a 12.2% rate among pregnant 20–24-year-olds. The prevalence of other sexually transmitted infections (STIs), combined with early sexual debut and low rates of condom use, facilitates the spread of HIV among Cameroonian youth (UNAIDS, 2002.) One study estimated condom use to be 16% among females aged 15–24 and 31% among males in the same age group (UNICEF, 2001). Cameroonian youth also experience high rates of unwanted pregnancies, abortion, and pregnancy-related school dropouts (Calvés & Meekers, 1997; Epanya & Delude, 1996; Kamtchouing et al., 1997).

In response to these growing threats to youth in Cameroon, a variety of reproductive health programs are being implemented throughout the country. One such program is *100% Jeune*, a social marketing reproductive health program targeting 15–24-year-olds living in Yaoundé and Douala, the two largest cities in Cameroon. The program uses a combination of mass media and interpersonal communication methods to motivate at-risk urban youth to use condoms consistently or not have sex. This study evaluates the *100% Jeune* program, analyzing its reach and impact on reproductive health outcomes, condom use, level of sexual activity, and predictors of condom use.

Background

The *100% Jeune* Adolescent Reproductive Health Program

With support from the Bill and Melinda Gates Foundation, the *100% Jeune* program was launched in Yaoundé and Douala in 2000 by Programme de Marketing Social au Cameroun (PMSC), an affiliate of Population Services International (PSI). *100% Jeune* aims to motivate at-risk urban youth to engage in healthier sexual behavior and focuses on reducing barriers to condom use. Program messages promote discussion of adolescent reproductive health issues within couples and between youth and their parents, highlight previous sexual history as a risk

factor for contracting STIs and HIV, and encourage young females to take responsibility for their reproductive health. The program promotes abstinence and consistent condom use as effective strategies to prevent unwanted pregnancy, STIs, and HIV. Mass media messages focus on increasing condom use with regular partners in particular.

The program uses a variety of mass media and interpersonal communication methods to diffuse its messages and encourage youth to practice and adopt new, healthy behaviors. Peer education sessions and a weekly radio call-in show are key interactive components of the program. Other important elements include a monthly magazine, *100% Jeune, Le Journal*, and a serial radio drama titled *Solange, Let's Talk about Sex*. Integrated television, radio, and billboard campaigns as well as a network of branded youth-friendly *Vendeurs des Amis des Jeunes* condom outlets support these activities. The *100% Jeune* program activities are integrated into a pre-existing national contraceptive social marketing program.

Messages and activities were refined in 2002 on the basis of mid-project research results and an external evaluation of the peer education program (Meekers et al., 2003; Wilson 2002). New program strategies implemented between the 2002 and 2003 survey waves include encouraging parents to discuss reproductive health issues with their children, making program materials and messages more appealing to females, and integrating participatory approaches into peer education sessions. Less emphasis was placed on promoting youth-friendly condom outlets during the final 18 months of the study period (Ward, 2003).

Theoretical Framework

100% Jeune is a theory-based program aiming to reduce barriers to condom use and increase safe behavior—namely abstinence or consistent condom use—among youth. Reproductive health programs often use popular theories such as Social Learning Theory, the Theory of Reasoned Action, and the Health Belief Model to identify behavior change program objectives and activities (Bandura, 1977; Becker, 1974; Janz & Becker, 1984; Mantell, DiVittis, & Auerbach, 1997; Rosenstock, Strecher, & Becker, 1988). The *100% Jeune* program is based on a comprehensive theoretical framework that uses key elements from all of these behavior change theories.

Recent studies suggest that condom use among youth is associated with several factors, including (1) perceived severity of sexual risks, (2) perceived personal risk (susceptibility), (3)

perceived condom attributes and access, (4) perceived social support, and (5) self-efficacy (Estrin, 1999; Lahia & Ross, 1997; MacPhail & Campbell, 2001; Maswanya et al., 1999; Meekers & Klein, 2002; Peltzer, 2000). This analysis measures progress and program impact in each of these five areas that predict condom use.

Methods

Data

This analysis uses data from three waves of the Cameroon Adolescent Reproductive Health Survey, implemented at 18-month intervals between 2000 and 2003. The surveys were commissioned by PMSC and implemented by IRESCO¹ in 2000 and 2002 and by FOCAP² in 2003. The first wave was implemented from July 26 to August 10, 2000, the second from January 4 to January 19, 2002 (Tchupo & Tégang, 2001; Tchupo & Tégang, 2002), and the third from June 19 to July 16, 2003 (FOCAP, 2003). The surveys contain information on randomly selected samples of 15–24-year-old youth living in the cities of Yaoundé and Douala. All surveys used the same base questionnaire that covered a range of reproductive health topics, including questions regarding HIV risk behavior, level of sexual activity, and condom use. The 2002 and 2003 surveys included additional questions to assess exposure to the *100% Jeune* program and other youth programs. The questionnaires for each of the three survey waves were pretested among a group of 30 male and female youth.

A four-stage stratified sampling design was used for all three surveys. In 2000, the target sample size was 2,000; this figure was increased to 3,500 in 2002 and 2003. The sample was equally divided between Yaoundé and Douala. Using probability of selection proportional to population size (PPS), neighborhoods (*quartiers*) were selected in each city. Twelve neighborhoods were selected for the 2000 survey and 20 were chosen for the 2002 and 2003 surveys. Again using PPS, a total of 30 enumeration areas were chosen in the selected neighborhoods. Households were selected randomly from a list of households in which at least one eligible person resided. One eligible person per household was randomly selected and up to

¹ L'Institut de Recherche et des Études de Comportements.

² Forum Camerounais de Psychologie.

three interview attempts were made. No replacements were made for respondents who could not be reached.

Same-sex interviewers, aged 25 or younger, participated in a three-day training workshop prior to conducting the interviews in 2000 and 2002, and interviewers in 2003 received four days of training because of the long length of the questionnaire. The fieldwork team for the 2000 survey consisted of 24 interviewers and 4 supervisors. The increased survey sample size in 2002 and 2003 required the size of the fieldwork team to be increased to 64 interviewers and 8 supervisors. A substantial number of data collectors worked on more than one of the three surveys. All interviews were conducted in French, the national language, and lasted approximately 45 minutes. Verbal informed consent was obtained from both the head of the household and the respondent. The final sample sizes for the three surveys were 2,096, 3,536, and 3,627, respectively. Our analysis is restricted to the subsample of unmarried³ youth, as this is the main target population for the program. This reduces the effective sample size to 1,956 youth in 2000, 3,237 in 2002, and 3,370 in 2003.

Measures

Program Exposure

To measure the effect of exposure to the *100% Jeune* program in 2002 and 2003, we created a variable based on the following program elements: (1) often or always listening to the *100% Jeune* radio call-in show, (2) having heard at least 10 episodes of the *Solange* radio drama, (3) often or always reading *100% Jeune, Le Journal*, and (4) having spoken to a *100% Jeune* peer educator. Respondents satisfying two or more of these criteria were classified as having “high” exposure, those who fulfilled one of the criteria as having “medium-high” exposure, and the remaining respondents as having “low” exposure.

Predictors of Condom Use

As previously described by Meekers et al. (2003), we analyze changes in variables shown to be associated with condom use. The predictor variables cover the following factors in the theoretical framework, described earlier, that was used to design the program. Although the program

³ “Unmarried” is defined as not currently married or living in union.

focuses on these factors to varying degrees, it is necessary to assess all of them to have a full understanding of changes in condom use and reproductive health.

- *Perceived severity of the health threat*: Respondents were asked, “Do you believe that a person who has HIV/AIDS can survive?” and “Do you believe that AIDS can be cured?” The indicators of perceptions of the health threat are two dichotomous variables indicating whether the respondent believes that someone infected with HIV/AIDS can survive and whether she or he believes that AIDS can be cured (“yes” vs. “no/don’t know”).
- *Perceived risk*: To measure perceived risk of HIV infection, respondents were asked, “If you would not use condoms, would you say your risk of contracting HIV/AIDS would be high, moderate, low, or there would be no risk?” The clause “if you would not use condoms” avoids the problem that some respondents may factor in condom use in their risk assessment while others may not. The variable measures whether respondents believe their risk to be moderate or high versus other. Youth were also asked if they know someone who has HIV/AIDS or who has died of HIV/AIDS (“yes” vs. “no/don’t know”).
- *Perceived condom attributes and access*: Respondents were asked whether condoms are effective for pregnancy prevention, for STI prevention, and for HIV/AIDS prevention. These two variables are coded as “yes” versus “no/don’t know.” We also asked youth, “Do you believe that condom use reduces sexual pleasure for men?” Our indicator is a dichotomous variable that equals one if the respondent perceives that condom use reduces sexual pleasure, zero otherwise. Finally, respondents were asked how long it would take to walk from their house to the nearest condom outlet. This information is coded as “knows condom source within 10 minutes” versus “does not know source within 10 minutes or does not know a source at all.”
- *Self-efficacy*: Because self-efficacy is known to be a multifaceted concept (Meekers & Klein, 2002; Parsons et al., 2000), several components of condom use self-efficacy were measured. For this study, respondents’ perceived ability to negotiate condom use, to obtain condoms, and to use them correctly are focused on. To measure confidence in their ability to negotiate condom use, interviewers asked respondents who had a regular partner in the past year, “Are you sure that you can convince him/her to use condoms with you (yes vs. no/not sure)?” Respondents were also asked, “Would you be shy buying condoms in a shop near your

home?” Our indicator is a dichotomous variable that equals one for respondents who indicated they would not be shy, zero for all others. Finally, to measure whether respondents believe that they have the skills to actually use a condom, respondents were asked, “Are you sure that you know how to correctly use a condom (“yes” vs. “no/not sure”)?”

— *Perceived social support:* Youth were asked several questions pertaining to perceived social support. They were asked if their parents support condom use by youth and if their friends are supportive of using condoms. Both questions are coded as “yes” versus “no/don’t know.” Youth were also asked if they discussed the use of family planning in the past year and whether they discussed STI/AIDS prevention in the past year (“yes” vs. “no/don’t know”).

Sexual Activity Indicators

Indicators of level of sexual activity include “had two or more sexual partners in the past year” and “had sex in the past year” (defined as having one or more sexual partners in the past year and measured among all respondents).

Condom Use Indicators

The question, “Have you ever used a condom?” was asked to measure the extent to which youth tried condoms. As youth who try condoms do not necessarily continue to use them, we also measured condom use in the last sexual act with a regular partner and use in the last sexual act with a casual partner. These indicators are coded as yes/no. Respondents were also asked about frequency of condom use with their regular and casual partners (never, sometimes, often, or always). These two variables measure consistent condom use and are recoded as “always used condoms” versus “often/sometimes/never used condoms.”

Reproductive Health Indicators

Sexually experienced males were asked if they had any STI symptoms in the past year. This variable is coded as “yes” versus “no/don’t know.” Sexually experienced females were asked about the timing of their last pregnancy—did they want to be pregnant then, to be pregnant later, or not have any more children? This variable measures unwanted pregnancy and is recoded as “would have preferred to wait/not have any more children” versus “wanted to be pregnant at that time.”

Control Variables

To account for confounding factors, control variables include age, city of residence, level of education, school enrollment status, socioeconomic status, number of sexual partners, and spontaneous recall of other adolescent reproductive health programs.

Methods

Logistic regression analyses examine trends in predictors of condom use as well as indicators of condom use and level of sexual activity. The results of the logistic regression models are presented as adjusted proportions. The trend analyses control for age, city of residence, level of education, school enrollment status, socioeconomic status, and number of sexual partners. All analyses are conducted separately for male and female respondents.

In a subsequent section, logistic regression analyses also study how predictors of condom use, level of sexual activity, condom use, and reproductive health outcomes vary with level of exposure to the *100% Jeune* program. These dose response analyses measure to what extent youth with medium and high program exposure in 2002 and 2003 have better results than those with low program exposure in 2002 (the reference group) after controlling for age, city of residence, level of education, school enrollment status, socioeconomic status, number of sexual partners, and spontaneous recall of other programs. Comparison of youth with low program exposure in 2002 and 2003 provides an estimate of secular trends not attributable to the program. These analyses are also estimated separately for male and female respondents.

Study Limitations

This study is subject to several limitations. As with all studies based on cross-sectional data, the direction of causal relationships cannot always be determined. However, the three survey waves provide information about the temporal order of events, which in turn can help establish causality. Also, behavioral outcomes are self-reported and therefore subject to reporting errors and biases. Finally, although information on spontaneous recall was collected for all programs, it was more difficult to obtain information about specific program elements for reproductive health programs other than *100% Jeune*. This difficulty limits the extent to which we can control for exposure to other programs.

Sample Description

Table 1 shows the characteristics of the study samples. For all three surveys, half of the youth reside in the city of Douala and half in Yaoundé, as specified by the study design. Results show that both the 2002 and 2003 survey populations are younger than respondents in 2000. However, the 2002 and 2003 survey populations do not differ with respect to age. While there are more males than females in each survey round, the gender distribution remains the same across surveys (approximately 55% male and 45% female). Results from the 2001 Demographic and Health Survey also show a higher percentage of males in the youth population. The 2000 and 2003 survey populations are similar in terms of the level of education attained—about 90% of youth reached at least a secondary school education—whereas the 2002 survey population is less well educated, with only 85% having a secondary school education. The three samples also differ in terms of the percentage of youth in school and socioeconomic status. These differences between the study populations are controlled for in the analysis.

Table 1 About Here

Results

Reach of *100% Jeune* and Other Youth Reproductive Health Programs

To measure unprompted exposure to the *100% Jeune* program and other youth reproductive health programs, the following approach was used. Respondents were reminded that reproductive health programs provide family planning, pregnancy, HIV/AIDS prevention, and

other services, and that they are implemented by either nongovernmental organizations or government agencies. Youth were then asked to cite all the youth reproductive health programs they have heard of. Results are shown in Appendix 1. The percentage of youth reporting having heard of at least one program increased significantly throughout the study period, from 35% in 2000 to 42% in 2002 and to 50% in 2003. Unprompted recall of *100% Jeune* increased dramatically from 1% in 2000 to 26% in 2002. Between 2002 and 2003, the level of unprompted recall remained constant at that level. No other program shows unprompted recall as high as that of *100% Jeune*. Only the *Cercle des Amis du Cameroun* (CERAC) program (19%), *Entre Nous Jeunes* (9%), and the program conducted by the Ministère de la Santé Publique (9%) had unprompted recall higher than 5% in 2003.

Prompted recall of specific *100% Jeune* program elements is shown in Table 2. Results are limited to the 2002 and 2003 survey rounds, as the 2000 survey occurred before the *100% Jeune* program was fully implemented. Table 2 shows that recall of several program elements was higher in 2003 than in 2002. Similarly, the percentage of youth reporting high levels of exposure to these program elements was higher at the time of the 2003 survey than in the 2002 survey. Approximately 12% of youth in 2002 and 2003 reported ever attending a peer education session, while the percentage who ever spoke to a peer educator increased slightly but significantly, from 8% to 11% between 2002 and 2003. Reach of the *100% Jeune*, *Le Journal* magazine, the *100% Jeune* radio call-in show, and the *Solange, Let's Talk about Sex* radio drama also increased. About three-quarters of all youth (74%) in 2002 reported having read the magazine at least once, compared with 90% in 2003. The level of exposure to the magazine also increased significantly between 2002 and 2003. The percentage of youth who report reading it often or always increased from 34% to 48%. The reach of the radio call-in show increased from 47% to 52% of youth between the last two surveys, and at the time of the 2003 survey, 32% of youth had listened to the *Solange, Let's Talk about Sex* radio drama at least once in the previous three months, compared with 26% in 2002.

In terms of exposure to mass media spots, about 70% of youth in both 2002 and 2003 reported seeing a television spot in the past three months, but exposure to radio spots decreased from 65% in 2002 to 59% in 2003. Exposure to youth-friendly condom vendors also decreased between 2002 and 2003. Only 20% of youth in 2003 report hearing of them in the past three

months, compared with 32% in 2002. The percentage of youth visiting a youth-friendly vendor also decreased from 9% to 5% between the two survey rounds.

Table 2 About Here

Trends in Condom Use, Level of Sexual Activity, and Predictors of Condom Use

Tables 3a and 3b illustrate trends in condom use, level of sexual activity, and predictors of condom use. The results of this logistic regression analysis are shown as adjusted percentages and have been controlled for age, city of residence, level of education, school enrollment status, socioeconomic status, and number of sexual partners.⁴ We tested the significance of trends observed between each survey wave (2000 vs. 2002, 2002 vs. 2003) as well as across the entire study period (2000 vs. 2003). Table 3a shows results for females and Table 3b shows results for males.

Among both males and females, almost all predictors of condom use improved during the study period, as did the indicators of condom use. However, levels of risky sexual activity did not decline. Among males, self-reported STI symptoms decreased, but among females, the frequency of unwanted pregnancies remained unchanged. Below, these findings are described in detail.

Tables 3a and 3b About Here

Perceived Severity: Youth's perception of the severity of HIV and AIDS decreased over the three-year period. Among both sexes, youth in 2003 were more likely than they were in 2000 to feel that an HIV-positive person can survive. Among females, the percentage believing that an HIV-positive person could survive increased from 14% in 2000 to 51% in 2003; among males, the percentage increased from 25% to 42%. Female youth in 2002 and 2003 were also more likely than female youth in 2000 to report that AIDS can be cured. These perceptions may reflect

⁴ Note that trends between 2000 and 2003 may differ slightly from those reported in earlier studies (Meekers et al., 2003). These differences stem from the fact that our analyses are based on a merged data set that contains all three survey waves, which implies that the mean values of the control variables differ somewhat from those used earlier.

an increased awareness of antiretroviral therapies that are prolonging the lives of those infected with HIV.

Perceived Risk: Trends in youth's perception of their perceived personal risk for HIV are mixed. Among females, the percentage reporting a moderate to high perceived risk for HIV infection increased from 52% in 2000 to 66% in 2002, but then declined to 58% in 2003. Among males, the corresponding percentage was constant at roughly 73% in 2000 and 2002, but then declined significantly to 59% in 2003.

Perceived Condom Attributes and Access: Youth's knowledge of where to obtain condoms and their opinions about them improved over the 36-month study period. Between 2002 and 2003, the percentage reporting knowledge of a condom source within a 10-minute walk increased from 62% to 74% among females, and from 80% to 90% among males. Although the vast majority of both female and male youth in 2000 reported that condoms are effective for family planning and effectively prevent HIV, positive trends are observed for both indicators between 2000 and 2003. The percentage of females stating that condoms are effective for family planning increased from 76% to over 85%; for males, it increased from 76% to 88%. Similar results are observed regarding the effectiveness of condoms for HIV prevention. Most of these improvements occurred between the 2002 and 2003 surveys. Youth of both sexes are also less likely to report that condoms reduce sexual pleasure in 2003 than in 2000.

Self-Efficacy: Results from the three surveys indicate that over time, Cameroonian youth generally became more comfortable buying condoms and negotiating condom use, and also gained confidence about correct condom use. The percentage of females who are not shy about buying condoms in a nearby store increased from 43% in 2000 to 57% in 2003; among males it increased from 61% to 78%. Although levels were high at baseline—over 80%—both females and males reported more frequently in 2003 than in previous years that they were able to convince their regular partners to use condoms. Confidence about knowing how to use condoms correctly also increased, from 38% to 53% among females and from 66% to 72% among males across the three survey waves.

Social Support: Social support for youth condom use increased. Youth are more likely to report that their friends support youth condom use (86% in 2003 vs. 74% in 2000 among females, and 89% vs. 83% among males). The percentage of youth stating that their parents support condom use also rose between 2000 and 2003. In addition, respondents more frequently discuss family planning as well as STI/AIDS in the past year.

Sexual Activity: No changes are noted across the three surveys in the percentage of youth who report having had sex in the past year—about 75% of female youth and 80% of males reported being sexually active in the past year. The percentage of female youth who reported having two or more sexual partners in the year before the survey increased from 15% in 2000 to 19% in 2002. However, between 2002 and 2003 the percentage decreased significantly from 19% to 13%. This latter percentage is not significantly different from the percentage observed at the 2000 baseline survey. Among males, no significant changes in the number of sexual partners were observed.

Condom Use: Tables 3a and 3b show that among both male and female youth, all measures of condom use improved steadily over time. Youth became more likely to report using a condom in their last sex act with a regular partner as well as with a casual partner. Specifically, the percentage of females who reported using a condom during their last intercourse with their regular partner increased from 33% in 2000 to 46% in 2002 and to 62% in 2003. For males, the corresponding percentages are 44%, 62%, and 76%, respectively. Similar increases were observed in consistent condom use, with both regular and casual partners. Consistent use with regular partners increased from 14% in 2000 to 21% in 2002 and to 31% in 2003 among females, and from 20% in 2000 to 33% in 2002 and to 45% in 2003 among males. Among females, consistent condom use with casual partners increased from 29% to 50% during the first phase of the project (between the 2000 and 2002 surveys) and from 50% to 70% during the second phase (between the 2002 and 2003 surveys). Among males, consistent use with casual partners increased from 45% to 53% during the first phase, and further increased to 70% by the time of the 2003 survey.

Reproductive Health: In terms of reproductive health outcomes, reported STIs declined over time but unwanted pregnancies did not. The percentage of sexually experienced males reporting an STI symptom in the past year decreased steadily, from 20% in 2000 to 11% in 2003.

The Effect of Exposure to the 100% *Jeune* Program

Tables 4a and 4b examine to what extent reproductive health outcomes, level of sexual activity, condom use, and predictors of condom use vary with level of exposure to the 100% *Jeune* program in 2002 and 2003. Because program activities were not fully implemented at the time of the 2000 survey, the analysis is limited to the latter two survey waves. To measure the impact of program exposure, we compared results among youth with medium and high exposure in each of the two survey waves (2002 and 2003) with those with low exposure in the 2002 survey, who serve as the reference group. To measure the extent of secular trends occurring independently of exposure to the 100% *Jeune* program, we compared youth with low exposure in the 2002 and 2003 surveys. The results of this logistic regression analysis are shown as adjusted percentages and have been controlled for age, city of residence, level of education, school enrollment status, socioeconomic status, number of sexual partners, and spontaneous recall of other reproductive health programs. Table 4a shows results among females and Table 4b reports results among males.

If a secular trend exists that cannot be attributed to the 100% *Jeune* program, we expect that even youth with low program exposure will show improvement between 2002 and 2003. If the 100% *Jeune* program contributed to observed improvement, we expect that youth with medium-high and/or high exposure in 2003 will have significantly better results for a variable than those in the reference group, and also that these results will be substantially better than any observed secular trend.

Tables 4a and 4b About Here

The results in Tables 4a and 4b indicate that improvement occurred for a variety of indicators, some of which can be attributed to the 100% *Jeune* program. Among both male and female youth, the 100% *Jeune* program contributed to increased use of condoms. Among females reporting high exposure to the 100% *Jeune* program in the 2003 survey, 39% report consistently

using condoms with regular partners, compared with only 20% among those in the 2002 low-exposure reference group. In contrast, the secular trend indicates improvement from 20% to only 27%. Among males, 53% of those with high exposure in 2003 report consistent condom use with their regular partners, compared with only 29% of those in the reference group. The secular trend among males shows improvement from 29% to only 38%. Among males, the program also contributed to improvements in condom use in the last sexual act with a casual partner and consistent use with a casual partner; 79% of male youth with high exposure in 2003 reported consistent condom use with a casual partner, compared with 48% among those in the reference group. In comparison, the secular trend indicated an improvement from only 48% to 63%. Condom use among females could not be assessed because of the small number of females reporting casual partners.

Between 2002 and 2003, the program also appears to have contributed to improvements in predictors of condom use, including perceived condom attributes, perceived social support, and self-efficacy among youth of both sexes. Among females, the data indicate that the program reduced shyness in purchasing condoms and increased discussion of family planning with people other than their friends. Among males, high program exposure in 2003 is associated with peer support of condom use, and no secular trend is noted. In addition, among both sexes, high program exposure is also associated with the belief that parents support condom use.

The program did not decrease the level of sexual activity or reduce the number of sexual partners, despite efforts to promote abstinence. Furthermore, it did not affect the percentage of males reporting STI symptoms or the frequency of females reporting unwanted pregnancies.

For several variables, there were observed secular trends not attributable to the *100% Jeune* program. This indicates that factors besides the *100% Jeune* program also contributed to improvements in outcome variables. For example, among females with low program exposure, the percentage who believe condoms are effective for family planning increased significantly, from 72% in 2002 to 84% in 2003. Among males, the percentage for this indicator increased from 82% to 86% between 2002 and 2003. Secular trends were also observed in the areas of perceived condom effectiveness, condom access, self-efficacy, social support, and condom use.

Conclusions

This paper examines the *100% Jeune* adolescent reproductive health program implemented in the cities of Douala and Yaoundé, Cameroon. The *100% Jeune* program targets 15–24-year-old youth using a combination of mass media and interpersonal communication channels to motivate them to use condoms consistently or not have sex. This study analyzes the program's reach and its impact on condom use, level of sexual activity, and predictors of condom use. The results are based on three rounds of an adolescent reproductive health survey conducted at 18-month intervals between 2000 and 2003.

Results show that *100% Jeune* successfully used a variety of mass media and interpersonal communication channels to reach a high proportion of youth throughout the intervention period. About one in four youth (26% and 28%, respectively) surveyed in both 2002 and 2003 spontaneously recalled the *100% Jeune* program. In a context in which a variety of governmental and nongovernmental partners are increasing youth-focused reproductive health programming, the *100% Jeune* program reached a higher proportion of youth than did other programs. In addition, the percentage of youth who recalled specific *100% Jeune* program elements was higher in the 2003 survey than in the 2002 survey. In the 2003 survey, youth were also more likely to report repeated exposure to campaign elements than was the case in 2002. Television spots and the monthly magazine reached the highest percentage of youth. Although only 12% of all youth were reached through peer education, the percentage of youth who reported speaking with a program peer educator was higher in 2003 than in the 2002 survey. This result may be related to the program's increased efforts to integrate participatory approaches into peer education sessions.

An examination of trends in behaviors and related predictors over the 36-month study period shows that substantial positive changes occurred among youth. Results of dose response analyses indicate that some of these positive changes can be attributed to the *100% Jeune* youth social marketing program. The program contributed to substantial increases in condom use, including consistent use with regular partners among youth of both sexes. Among males, it also contributed to consistent use with casual partners. Whereas the program had some impact on condom use among males after the first 18 months of activities, the 2003 results reflect increased consistent condom use among both females and males with high exposure to the *100% Jeune* program. Results also indicate that the *100% Jeune* program contributed to important

improvements in predictors of condom use. The program did not decrease the level of sexual activity or reduce the number of sexual partners, despite efforts to promote abstinence. Furthermore, it did not affect the percentage of males reporting STI symptoms or females reporting unwanted pregnancies.

In addition, the observed secular trends indicate that factors besides *100% Jeune* also contributed to the improvement observed in the areas of perceived condom effectiveness, condom access, self-efficacy, social support, and condom use. These results indicate that collective efforts of multiple organizations over time can lead to improvements in adolescent reproductive health.

Although there are signs of increasing use of condoms and reduced barriers among Cameroonian youth, several challenges remain. Youth of both sexes continue to believe they are not at risk of contracting HIV/AIDS, and social barriers continue to limit open discussion of reproductive health issues. Females still appear to be less likely than males to believe they can buy and use condoms correctly. Future programs should aim to address these and other barriers to improved adolescent reproductive health. Programs should devote resources to motivating youth to abstain, or reduce numbers of sexual partners, and continue to encourage sexually active youth to use condoms consistently, that is, with every partner, every time. Resources should be allocated to identify and understand predictors of abstinence and partner reduction to inform future programming decisions.

Competing Interests

Miss Plautz is Research Associate with Population Services International (PSI), which is the mother company of the organization that implemented the program that is being assessed in this study. Dr. Meekers is the former Research Director of PSI, and was a paid consultant on this project.

Author's Contributions

Miss Plautz conducted the statistical analyses and prepared the first draft of this manuscript. Dr. Meekers developed the study design, assisted with the analysis, and contributed to the writing of the final manuscript.

Acknowledgments

This paper was funded by Population Services International (PSI) through a grant for adolescent reproductive health social marketing programs from the Bill and Melinda Gates Foundation. The authors are grateful to Jim Foreit of the Population Council and Josselyn Neukom of PSI for commenting on an earlier version of this paper and to Catherine Malo at EEI Communications for editing it.

References

- Bandura, A. 1977. *Social Learning Theory*. Englewood Cliffs, NJ: Prentice-Hall.
- Becker, M. 1974. The Health Belief Model and personal health behavior. *Health Education Monographs*, 2, 236–473.
- Calvés, A. E., & Meekers, D. 1997. *Gender Differentials in Premarital Sex, Condom Use, and Abortion: A Case Study of Yaoundé, Cameroon*. Washington, DC: Population Services International.
- Epanya, A., & Delude, B. R. 1996. Country Watch: Cameroon. *AIDS/STD Health Promotion Exchange*, 3, 9–10.
- Estrin, D. 1999. In Ghana, young men's condom use is linked to lack of barriers, perceived susceptibility to HIV infection. *International Family Planning Perspectives*, 25(2), 106–111.
- Forum Camerounais de Psychologie. 2003. Enquête sur la santé reproductive des adolescents au Cameroun (Yaoundé et Douala), 2003. Unpublished report.
- Janz, N. K., & Becker, M. H. 1984. The Health Belief Model: A decade later. *Health Education Quarterly*, 11(1), 1–47.
- Kamtchouing, P., Takougang, I., Ngoh, N., & Yakam, I. 1997. La sexualite des adolescents en milieu scolaire a Yaounde (Cameroun). *Contraception, Fertility, Sexuality*, 25(10), 798–801.
- Lahia, M., & Ross, M. 1997. HIV/AIDS prevention-related social skills and knowledge among adolescents in Sierra Leone, West Africa. *African Journal of Reproductive Health*, 1(1), 37–44.
- MacPhail, C., & Campbell, C. 2001. 'I think condoms are good but, aai, I hate those things': Condom use among adolescents and young people in a Southern African township. *Social Science & Medicine*, 52(11), 1613–1628.
- Mantell, J. E., DiVittis, A. T., & Auerbach, M. I. 1997. Applying theory to HIV prevention interventions. In J. D. A. Mantell & M. Auerbach (Eds.), *Evaluating HIV Prevention Interventions* (pp. 179–204). New York: Plenum.
- Maswanya, E., Moji, K., Horiguchi, I., Nagata, K., Aoyagi, K. I., & Honda, S. T. T. 1999. Knowledge, risk perception of AIDS and reported sexual behavior among students in secondary schools and colleges in Tanzania. *Health Education Research*, 14(2), 185–196.
- Matagne, A. K., Tapko, J. B., Kamdem, T. K., Zokong, B. Z., Tookam, M. T., & Kaptué, L. N. 1998. Evolution of HIV Seroprevalence in the General Hospital Yaoundé. Paper presented at the International Conference on AIDS, Geneva, 1998.

- Mbopi, F., Mbu, R., Mauclere, P., Andela, A., Tetanye, E., Leke, R., Chaout, G., Barre Sinoussi, F., Martin, P., & Belec, L. 1998. Antenatal HIV prevalence in Yaoundé, Cameroon. *International Journal of STD and AIDS*, 9(7), 400–402.
- Meekers, Dominique, Sohail Agha and Megan Klein. 2003. The Impact on Condom Use of the “100% Jeune” Social Marketing Program in Cameroon. PSI Research Division, Working Paper No. 54. Washington, DC: Population Services International.
- Meekers, Dominique & Megan Klein. 2002. Understanding gender differences in condom use self-efficacy among youth in urban Cameroon. *AIDS Education and Prevention*, 14(1), 62–72.
- Parsons, J., Halkitis, P., Bimbi, D., & Borkowski, T. 2000. Perceptions of the benefits and costs associated with condom use and unprotected sex among late adolescent college students. *Journal of Adolescence*, 23(4), 377–391.
- Peltzer, K. 2000. Factors affecting condom use among senior secondary school pupils in South Africa. *Central African Journal of Medicine*, 46(11), 302–308.
- Rosenstock, I. M., Strecher, V. J., & Becker, M. H. 1988. Social Learning Theory and the Health Belief Model. *Health Education Quarterly*, 15(2), 175–183.
- Tchupo, J. P., & Tégang, S. P. 2001. Enquête sur la santé reproductive des adolescents au Cameroun (Yaoundé et Douala), 2000. Washington, DC: Population Services International, IRESCO.
- Tchupo, J. P., & Tégang, S. P. 2002. Enquête sur la santé reproductive des adolescents au Cameroun (Yaoundé et Douala), 2002. Washington, DC: Population Services International, IRESCO.
- UNAIDS. 2002. Epidemiological Fact Sheets on HIV/AIDS and Sexually Transmitted Infections: Cameroon. 2002 Update. www.unaids.org [accessed September 10, 2003].
- UNICEF. 2001. Enquête á Indicateurs Multiples (MICS) au Cameroun 2000. Ministère de l’Economie et des Finances, Gouvernement du Cameroun.
- UNICEF. 2002. *The State of the World's Children 2002*. New York: UNICEF.
- Ward, Dana. 2003. PSI/Programme de Marketing Social au Cameroun. Personal communication.
- Wilson, David. 2002, April 25. PSI Youth Sexual Health Programme: Cameroon, Rwanda, Madagascar. Unpublished report.

Table 1: Sample Characteristics

	Survey Wave			Significance		
	2000	2002	2003	2000 vs. 2002	2002 vs. 2003	2000 vs. 2003
Age				***		**
15–19	57.3	61.1	60.3			
20–24	42.7	38.9	39.7			
Gender						
Male	54.0	54.2	55.0			
Female	46.0	45.8	45.0			
Education				***	***	
Secondary +	90.1	84.7	91.4			
< Secondary	9.9	15.3	8.6			
Student				***	***	***
No	33.6	42.7	27.8			
Yes	66.4	57.3	72.2			
Socioeconomic Status				***	***	***
Low	27.6	31.6	28.7			
Medium	38.1	42.5	45.3			
High	34.3	25.9	26.1			
City						
Yaoundé	50.9	48.9	49.3			
Douala	49.1	51.1	50.7			
Number of Cases	1,956 (100%)	3,237 (100%)	3,370 (100%)			

Note: *** p < .01; ** p < .05.

Table 2: Percentage of Youth Who Recall Exposure to Various *100% Jeune* Campaign Elements (prompted recall, 2002 and 2003 survey waves only)

Campaign Element	Survey Wave		Significance
	2002 (%)	2003 (%)	
Peer Education and Promotion			
Ever attended peer education session	11.9	12.3	
Ever talked to peer educators	7.6	10.6	***
<i>100% Jeune, Le Journal Magazine</i>			
Ever read <i>100% Jeune</i> magazine	73.9	89.8	***
How often do you read the magazine?			***
- Never	26.2	10.2	
- Sometimes	39.7	42.3	
- Often	18.3	28.4	
- Always	15.9	19.1	
<i>100% Jeune Radio Call-In Show</i>			
Heard <i>100% Jeune</i> radio program in past 3 months	47.3	51.5	***
How often do you usually listen?			***
- Never	52.7	48.5	
- Sometimes	31.1	34.9	
- Often	11.2	13.7	
- Always	5.1	2.8	
<i>Solange, Let's Talk about Sex Radio Drama</i>			
Heard <i>Solange</i> radio series in past 3 months	26.0	31.5	***
How many different episodes have you heard?			***
- None	74.0	68.5	
- One to four	13.9	18.9	
- Five to nine	6.3	8.2	
- Ten or more	5.8	4.4	
Mass Media Campaign			
Saw TV spot in past 3 months	70.7	69.9	
Heard radio spot in past 3 months	64.5	58.9	***
Youth-Friendly Vendors			
Heard about Vendeurs Amis des Jeunes outlets in past 3 months	31.9	19.7	***
Visited a Vendeurs Amis des Jeunes outlet in past 3 months	8.8	4.5	***
<hr/>			
Number of Cases	3,237	3,370	

Note: *** p < .01.

Table 3a: Logistic Regression Results of Trends in Predictors of Condom Use and Trends in Indicators of Condom Use, Adjusted Percentages among Females

	Females			Significance		
	2000	2002	2003	2000 vs. 2002	2002 vs. 2003	2000 vs. 2003
Perceived Severity						
-HIV-positive person can survive	14.3	35.5	50.9	***	***	***
-AIDS can be cured	10.9	17.2	17.2	***		***
Perceived Risk						
-Moderate/high personal risk	51.7	65.7	57.5	***	***	***
-Knows someone who has/died of HIV/AIDS	45.0	45.8	47.6			
Perceived Condom Attributes and Access						
-Condoms effective for FP	75.5	72.5	85.2		***	***
-Condoms effective for HIV/AIDS prevention	89.8	88.7	93.6		***	***
-Condom source within 10 minutes	62.1	68.1	74.0	***	***	***
-Condoms reduce sexual pleasure	36.0	39.1	30.1		***	***
Self-Efficacy						
- Can convince regular partner to use condoms ^a	80.5	83.0	89.9		***	***
- Can convince casual partner to use condoms ^b	86.1	83.3	92.4		**	
- Not shy to obtain condoms in nearby shop	42.7	55.3	57.4	***		***
- Confident knows correct condom use	38.3	46.4	53.1	***	***	***
Perceived Social Support						
-Friends support youth condom use	74.0	81.1	86.4	***	***	***
-Parents support youth condom use	58.4	68.0	71.3	***		***
-Discussed family planning with friends in past year	17.8	21.8	29.4		***	***
-Discussed family planning with others in past year	28.0	25.4	33.8		***	**
-Discussed STI/AIDS with friends in past year	32.5	39.2	44.3	**	**	***
-Discussed STI/AIDS with others in past year	38.6	31.7	45.2	***	***	**
Sexual Activity						
-Had sex in the past year	71.5	74.1	74.4			
-Had 2+ sexual partners in the past year	15.1	19.3	13.2	***	***	
Condom Use						
-Ever used condoms	50.0	61.6	64.9	***		***
-Used condom in last sex with regular partner ^a	32.9	46.0	62.1	***	***	***
-Always uses condoms with regular partner ^a	13.7	21.0	31.4	***	***	***
-Used condom in last sex with casual partner ^b	46.1	65.5	84.1	**	***	***
-Always uses condoms with casual partner ^b	29.1	49.6	69.9	***	***	***
Reproductive Health						
-Last pregnancy was unwanted ^c	71.0	68.6	58.6			
Number of Cases (100%)	899	1,483	1,518			

Note: *** p < .01; ** p < .05. FP = family planning; HIV = human immunodeficiency virus; STI/AIDS = sexually transmitted infections/acquired immunodeficiency syndrome.

Analysis controls for age, city of residence, level of education, school enrollment, socioeconomic status, and number of sexual partners.

Trends between 2000 and 2003 may differ slightly from those reported in earlier studies (Meekers et al., 2003). These differences stem from the fact that our analyses are based on a merged data set containing all three survey waves, which implies that the mean values of the control variables differ somewhat from those used earlier.

^a N = 474, 902, and 848, respectively.

^b N = 76, 105, and 115, respectively.

^c Among women reporting having had at least one pregnancy. N = 210, 55, and 39, respectively.

Table 3b: Logistic Regression Results of Trends in Predictors of Condom Use and Trends in Indicators of Condom Use, Adjusted Percentages among Males

	Males			Significance		
	2000	2002	2003	2000 vs. 2002	2002 vs. 2003	2000 vs. 2003
Perceived Severity						
-HIV-positive person can survive	25.2	38.9	42.2	***	**	***
-AIDS can be cured	21.1	22.6	21.6			
Perceived Risk						
-Moderate/high personal risk	73.4	73.8	59.3		**	***
-Knows someone who has/died of HIV/AIDS	36.5	32.6	39.7	**	***	
Perceived Condom Attributes and Access						
-Condoms effective for FP	79.4	82.9	87.5	**	***	***
-Condoms effective for HIV/AIDS prevention	88.0	87.8	90.8		***	**
-Condom source within 10 minutes	80.0	84.1	90.4	***	***	***
-Condoms reduce sexual pleasure	55.5	54.1	45.0		***	***
Self-Efficacy						
- Can convince regular partner to use condoms ^a	86.8	89.9	92.9		**	***
- Can convince casual partner to use condoms ^b	89.0	90.1	93.8		**	**
- Not shy to obtain condoms in nearby shop	61.1	68.1	78.1	***	***	***
- Confident knows correct condom use	65.9	71.9	71.9	***		***
Perceived Social Support						
-Friends support youth condom use	83.0	85.0	88.7		***	***
-Parents support youth condom use	64.3	74.8	75.6	***		***
-Discussed family planning with friends in past year	27.7	28.5	36.9		***	***
-Discussed family planning with others in past year	37.1	26.2	35.9	***	***	
-Discussed STI/AIDS with friends in past year	49.4	59.0	62.8	***		***
-Discussed STI/AIDS with others in past year	46.7	36.8	49.3	***	***	
Sexual Activity						
-Had sex in the past year	80.4	80.8	82.4			
-Had 2+ sexual partners in the past year	31.9	35.3	32.7			
Condom Use						
-Ever used condoms	58.7	64.9	70.2	**	***	***
-Used condom in last sex with regular partner ^a	44.3	61.5	75.7	***	***	***
-Always uses condoms with regular partner ^a	19.5	33.2	44.8	***	***	***
-Used condom in last sex with casual partner ^b	61.0	68.7	81.3	**	***	***
-Always uses condoms with casual partner ^b	45.3	52.5	70.1		***	***
Reproductive Health						
-Had an STI symptom in the past year ^c	19.6	14.7	10.7	***	***	***
Number of Cases (100%)	1,057	1,754	1,852			

Note: *** p < .01; ** p < .05. FP = family planning; HIV = human immunodeficiency virus; STI/AIDS = sexually transmitted infections/acquired immunodeficiency syndrome.

Analysis controls for age, city of residence, level of education, school enrollment, socioeconomic status, and number of sexual partners.

Trends between 2000 and 2003 may differ slightly from those reported in earlier studies (Meekers et al., 2003). These differences stem from the fact that our analyses are based on a merged data set containing all three survey waves, which implies that the mean values of the control variables differ somewhat from those used earlier.

^a N = 541, 978, and 974, respectively.

^b N = 282, 464, and 456, respectively.

^c Among sexually experienced males only because STIs are often asymptomatic among females. N = 721, 1,247, and 1,220, respectively.

Table 4a: Logistic Regression Results of Level of Exposure to the 100% Jeune program on Outcome Measures, Adjusted Percentages among Females

	2002			2003		
	Low exposure (ref.)	Medium-high exposure	High exposure	Low exposure	Medium-high exposure	High exposure
Perceived Severity						
-HIV-positive person can survive	32.7	38.1	37.7	48.2***	53.6***	51.5***
-AIDS can be cured	17.5	17.6	14.0	18.8	16.0	15.8
Perceived Risk						
-Moderate/high personal risk	64.0	65.7	70.9	55.6***	59.6	56.7**
-Knows someone who has/died of HIV/AIDS	41.0	50.0***	52.0***	45.9	50.4***	44.0
Perceived Condom Attributes and Access						
-Condoms effective for FP	72.4	72.4	73.3	84.1***	84.6***	89.0***
-Condoms effective for HIV/AIDS prevention	88.4	88.4	60.6	93.6***	94.2***	93.7**
-Condom source within 10 minutes	65.0	69.2	74.4**	72.2***	77.2***	71.9**
-Condoms reduce sexual pleasure	40.5	39.0	31.9**	33.6***	30.1***	23.3***
Self-Efficacy						
- Can convince regular partner to use condoms ^a	82.4	82.7	86.9	89.0***	90.8***	90.6**
- Can convince casual partner to use condoms ^b	--	--	--	--	--	--
- Not shy to obtain condoms in nearby shop	51.9	55.6	65.8***	51.9	60.0***	65.0***
- Confident knows correct condom use	38.1	50.1***	64.5***	47.6***	55.5***	59.4***
Perceived Social Support						
-Friends support youth condom use	78.3	82.3	85.8**	84.5***	87.2***	88.1***
-Parents support youth condom use	66.6	68.7	69.0	67.6	74.8***	71.1
-Discussed family planning with friends in past year	16.8	27.5***	25.3**	27.6***	24.6***	38.7***
-Discussed family planning with others in past year	23.4	23.9	33.0**	28.6	31.2**	47.8***
-Discussed STI/AIDS with friends in past year	35.2	43.3**	46.9**	42.0**	43.0**	49.1***
-Discussed STI/AIDS with others in past year	28.8	31.6	39.8**	41.7***	41.6***	58.9***
Sexual Activity						
-Had sex in the past year	77.8	75.7	77.2	75.5	77.5	80.4
-Had 2+ sexual partners in the past year	17.7	22.0	19.3	13.8**	12.8**	12.8
Condom Use						
-Ever used condoms	61.4	62.2	67.3	64.2	67.9**	67.5
-Used condom in last sex with regular partner ^a	47.3	42.2	50.0***	60.8***	60.0***	68.5***
-Always uses condoms with regular partner ^a	20.2	20.8	24.8	26.7**	32.6***	39.2***
-Used condom in last sex with casual partner ^b	--	--	--	--	--	--
-Always uses condoms with casual partner ^b	--	--	--	--	--	--
Reproductive Health						
-Last pregnancy was unwanted ^b	--	--	--	--	--	--
Number of Cases (100%)	789	494	200	640	589	289

Note: *** p < .01; ** p < .05. FP = family planning; HIV = human immunodeficiency virus; STI/AIDS = sexually transmitted infections/acquired immunodeficiency syndrome.

Analysis controls for age, city of residence, level of education, school enrollment status, socioeconomic status, number of sexual partners, and spontaneous recall of other programs.

^a N = 519, 275, and 108 in 2002 and 371, 314, and 163 in 2003.

^b Results are omitted because of the small number of cases.

Table 4b: Logistic Regression Results of Level of Exposure to the 100% *Jeune* program on Outcome Measures, Adjusted Percentages among Males

	2002			2003		
	Low exposure (ref.)	Medium-high exposure	High exposure	Low exposure	Medium-high exposure	High exposure
Perceived Severity						
-HIV-positive person can survive	39.6	37.9	37.5	42.1	42.3	42.0
-AIDS can be cured	23.7	21.2	20.9	22.0	21.2	21.5
Perceived Risk						
-Moderate/high personal risk	71.4	77.7**	78.7**	55.7***	63.6***	61.5***
-Knows someone who has/died of HIV/AIDS	30.0	35.9**	35.3	35.6**	44.9***	42.0***
Perceived Condom Attributes and Access						
-Condoms effective for FP	81.8	83.0	86.1	86.1**	88.7***	89.8***
-Condoms effective for HIV/AIDS prevention	86.6	90.0	88.4	90.4***	92.2***	89.8
-Condom source within 10 minutes	82.4	86.9**	83.8	89.7***	92.1***	88.9***
-Condoms reduce sexual pleasure	53.1	57.5	51.4	49.2	43.7***	35.9***
Self-Efficacy						
- Can convince regular partner to use condoms ^a	87.9	92.5**	93.9**	92.1**	93.8***	93.5
- Can convince casual partner to use condoms ^b	87.7	91.3	97.0**	93.8**	92.7	95.5
- Not shy to obtain condoms in nearby shop	64.8	73.8***	70.0	75.3***	82.1***	79.3***
- Confident knows correct condom use	67.9	76.8***	75.9**	66.3	77.5***	77.2***
Perceived Social Support						
-Friends support youth condom use	83.2	87.4**	89.0**	86.3	91.5***	90.5***
-Parents support youth condom use	73.0	76.5	77.1	71.0	80.9***	78.9**
-Discussed family planning with friends in past year	25.5	29.5	36.1***	39.1***	32.0**	36.5***
-Discussed family planning with others in past year	22.5	28.8**	32.8***	34.7***	32.6***	40.5***
-Discussed STI/AIDS with friends in past year	54.6	64.8***	67.1***	63.5***	60.9**	63.0**
-Discussed STI/AIDS with others in past year	31.8	38.6**	49.7***	44.1***	50.7***	58.6***
Sexual Activity						
-Had sex in the past year	79.4	81.5	84.3	81.7	83.1	83.9
-Had 2+ sexual partners in the past year	33.8	37.2	36.2	32.7	30.5	35.7
Condom Use						
-Ever used condoms	60.9	71.7***	69.6**	68.2**	73.9***	75.9***
-Used condom in last sex with regular partner ^a	57.7	62.8	73.0***	70.2***	80.8***	81.6***
-Always uses condoms with regular partner ^a	29.4	35.1	39.5**	37.7***	52.1***	52.7***
-Used condom in last sex with casual partner ^b	66.1	76.2**	65.5	77.8***	85.6***	89.3***
-Always uses condoms with casual partner ^b	48.0	64.7***	45.1	63.4***	79.9***	79.3***
Reproductive Health						
-Had an STI symptom in the past year ^c	16.1	12.8	12.0	11.2***	8.4***	11.6
Number of Cases (100%)	976	504	274	929	601	322

Note: *** p < .01; ** p < .05. FP = family planning; HIV = human immunodeficiency virus; STI/AIDS = sexually transmitted infections/acquired immunodeficiency syndrome.

Analysis controls for age, city of residence, level of education, school enrollment status, socioeconomic status, number of sexual partners, and spontaneous recall of other programs.

^a N = 532, 282, and 164 in 2002 and 483, 314, and 177 in 2003.

^b N = 261, 138, and 65 in 2002 and 249, 139, and 68 in 2003.

^c Among sexually experienced males only because STIs are often asymptomatic among females. N = 686, 367, and 194 in 2002 and 624, 389, and 207 in 2003.

Appendix 1: Percentage of Youth Aware of Specific Reproductive Health Programs, by Survey Wave (spontaneous responses only)

	Survey Wave			Significance		
	2000	2002	2003	2000 vs. 2002	2002 vs. 2003	2000 vs. 2003
Cameroon National Association for Family Welfare (CAMNAFAW)	2.2	1.4	2.1		**	
Femme-Santé Développement (FESADE)	1.9	0.6	0.9	***		***
Entre Nous Jeunes	4.0	3.5	9.1		***	***
Youth Development Association (YDF)	1.1	0.3	1.2	***	***	
Education à la Vie et l'Amour (SCS/EVA)	0.8	0.5	0.3			
SCOUTS	0.4	0.3	1.1		***	***
Croix Rouge	2.2	1.3	3.4		***	
EEC Appel	0.3	0.4	1.5		***	***
100% Jeune (PMSC)	1.3	25.9	27.9	***		***
Ministère de l'Education Nationale	2.5	0.9	2.5	***	***	
Ministère de la Santé Publique	12.4	4.4	8.9	***	***	***
Ministère de la Jeunesse et des Sports	0.6	0.3	1.9		***	***
Cercle des Amis du Cameroun (CERAC)	^a	14.8	19.4		***	
Other Programs	14.8	6.8 ^b	8.9 ^b			
Any Program	34.8	41.9	49.8	***	***	**
Number of Cases	1,956	3,236	3,370			

Note: *** p < .01; ** p < .05.

^a Not coded in 2000 survey wave

^b Significance not tested because slight changes in definition of "other."